

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Mammaglobin (SCGB2A2) (Breast Cancer Marker) (MGB1/2123R), CF640R conjugate, 0.1mg/mL, Clone: [MGB1/2123R], Rabbit, Monoclonal BOT-BNC402123-500

| | |
|--------------------------|---|
| Artikelname | Mammaglobin (SCGB2A2) (Breast Cancer Marker) (MGB1/2123R), CF640R conjugate, 0.1mg/mL, Clone: [MGB1/2123R], Rabbit, Monoclonal |
| Artikelnummer | BOT-BNC402123-500 |
| Hersteller Artikelnummer | BNC402123-500 |
| Alternativnummer | BOT-BNC402123-500-500UL |
| Hersteller | Biotium |
| Wirt | Rabbit |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant full-length human Mammaglobin (SCGB2A2) protein |
| Konjugation | CF640R |
| Produktbeschreibung | Mammaglobin is a 93 amino acid glycoprotein with homology to other secretoglobin-uteroglobin family members. It was originally identified as a breast cancer restricted biomarker by differential screening. Mammaglobin related to secretoglobin family t... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [MGB1/2123R] |
| Molekulargewicht | 18 kDa |
| UniProt | Q13296 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.1-0.2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |