

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2151), CF640R conjugate, 0.1mg/mL, Clone: [TACSTD2/2151], Mouse, Monoclonal BOT-BNC402151-100

| | |
|--------------------------|---|
| Artikelname | TACSTD2 / TROP2 (Epithelial Marker) (TACSTD2/2151), CF640R conjugate, 0.1mg/mL, Clone: [TACSTD2/2151], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC402151-100 |
| Hersteller Artikelnummer | BNC402151-100 |
| Alternativnummer | BOT-BNC402151-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary) |
| Konjugation | CF640R |
| Produktbeschreibung | TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites ne... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [TACSTD2/2151] |
| Molekulargewicht | 40 kDa |
| UniProt | P09758 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |