

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Emerin (Papillary Thyroid Carcinoma and EDMD Marker) (EMD/2167), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [EMD/2167], Mouse, Monoclonal BOT-BNC402167-500**

Artikelname	Emerin (Papillary Thyroid Carcinoma and EDMD Marker) (EMD/2167), CF640R conjugate, 0.1mg/mL, IgG1, Clone: [EMD/2167], Mouse, Monoclonal
Artikelnummer	BOT-BNC402167-500
Hersteller Artikelnummer	BNC402167-500
Alternativnummer	BOT-BNC402167-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, IHC, WB
Spezies Reaktivität	Human
Immunogen	Recombinant human Emerin protein fragment (around aa 56-167) (exact sequence is proprietary)
Konjugation	CF640R
Produktbeschreibung	Emerin is a member of the nuclear lamina associated protein family. It is ubiquitously expressed and localized to the nuclear membrane in normal cells. Mutations of the gene that encodes emerlin result in the X-linked recessive disease Emery-Dreifuss ...
Klonalität	Monoclonal

Konzentration	0.1 mg/mL
Klon-Bezeichnung	[EMD/2167]
Molekulargewicht	37 kDa
Isotyp	IgG1
UniProt	<a href="#">P50402</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user