

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 8 (KRT8) (KRT8/2174R), CF640R conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal BOT-BNC402174-100

Artikelname	Cytokeratin 8 (KRT8) (KRT8/2174R), CF640R conjugate, 0.1mg/mL, Clone: [KRT8/2174R], Rabbit, Monoclonal
Artikelnummer	BOT-BNC402174-100
Hersteller Artikelnummer	BNC402174-100
Alternativnummer	BOT-BNC402174-100-100UL
Hersteller	Biotium
Wirt	Rabbit
Kategorie	Antikörper
Applikation	IHC, WB
Spezies Reaktivität	Bovine, Human, Porcine, Rabbit, Sheep
Immunogen	Recombinant full-length human KRT8 protein
Konjugation	CF640R
Produktbeschreibung	Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[KRT8/2174R]

Molekulargewicht	52.5 kDa
UniProt	P05787
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin): 0.5-1.0 ug/mL for 30 minutes at RT Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user