

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

FOXP3(FXP3/197), CF647 conjugate, 0.1mg/mL, Clone: [FXP3/197], Mouse, Monoclonal BOT-BNC470197-100

| | |
|--------------------------|---|
| Artikelname | FOXP3(FXP3/197), CF647 conjugate, 0.1mg/mL, Clone: [FXP3/197], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470197-100 |
| Hersteller Artikelnummer | BNC470197-100 |
| Alternativnummer | BOT-BNC470197-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Monkey, Mouse |
| Immunogen | Recombinant human full-length FOXP3 protein |
| Konjugation | CF647 |
| Produktbeschreibung | Recognizes a protein of 47-55 kDa, which is identified as FOXP3. Its precise epitope is not known, but it has been mapped to the N-terminal portion of the protein. The FOX family of transcription factors is a large group of proteins that share a comm... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [FXP3/197] |

| | |
|------------------------|--|
| Molekulargewicht | 47-55 kDa |
| UniProt | Q9BZS1 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |