

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MHC II (HLA-DP) (beta chain)(BRA-FB6), CF647 conjugate, 0.1mg/mL, Clone: [BRA-FB6], Mouse, Monoclonal BOT-BNC470199-100**

|                          |  |
|--------------------------|--|
| Artikelname              | MHC II (HLA-DP) (beta chain)(BRA-FB6), CF647 conjugate, 0.1mg/mL,<br>Clone: [BRA-FB6], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC470199-100  |
| Hersteller Artikelnummer | BNC470199-100  |
| Alternativnummer         | BOT-BNC470199-100-100UL  |
| Hersteller               | Biotium  |
| Wirt                     | Mouse  |
| Kategorie                | Antikörper   |
| Applikation              | ELISA, FC, Functional Studies, IP  |
| Spezies Reaktivität      | Human  |
| Immunogen                | Non-T, non-B human acute lymphoblastic leukemia REH6 cell line   |
| Konjugation              | CF647  |
| Produktbeschreibung      | Recognizes a non-polymorphic determinant of DP-MHC class II. MHC class II antigens are transmembrane glycoproteins of non-covalently linked alpha (33-35 kDa) and beta (27-30 kDa) chains. It reportedly reacts with B- & non-T, non-B cell lines but not w... |
| Klonalität               | Monoclonal   |
| Konzentration            | 0.1 mg/mL  |
| Klon-Bezeichnung         | [BRA-FB6]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 33-35 kDa  |
| UniProt                | <a href="#">P04440</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |