

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD2(LFA2/600), CF647 conjugate, 0.1mg/mL, Clone: [LFA2/600], Mouse, Monoclonal BOT-BNC470600-100**

|                          |   |
|--------------------------|---|
| Artikelname              | CD2(LFA2/600), CF647 conjugate, 0.1mg/mL, Clone: [LFA2/600], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC470600-100   |
| Hersteller Artikelnummer | BNC470600-100   |
| Alternativnummer         | BOT-BNC470600-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF  |
| Spezies Reaktivität      | Human, Porcine, Primate   |
| Immunogen                | Recombinant human LFA2 protein  |
| Konjugation              | CF647   |
| Produktbeschreibung      | CD2 interacts through its amino-terminal domain with the extracellular domain of CD58 (also designated CD2 ligand) to mediate cell adhesion. CD2/CD58 binding can enhance antigen-specific T cell activation. CD2 is a transmembrane glycoprotein that is ... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [LFA2/600]  |
| Molekulargewicht       | ~50 kDa   |
| UniProt                | <a href="#">P06729</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |