

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD66 (CEA)(CEA31), CF647 conjugate, 0.1mg/mL, Clone: [CEA31], Mouse, Monoclonal BOT-BNC470670-500

| | |
|--------------------------|---|
| Artikelname | CD66 (CEA)(CEA31), CF647 conjugate, 0.1mg/mL, Clone: [CEA31], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470670-500 |
| Hersteller Artikelnummer | BNC470670-500 |
| Alternativnummer | BOT-BNC470670-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Monkey |
| Immunogen | Human colon carcinoma extract |
| Konjugation | CF647 |
| Produktbeschreibung | This antibody recognizes proteins of 80-200 kDa, identified as different members of CEA family. CEA is synthesized during development in the fetal gut and is re-expressed in increased amounts in intestinal carcinomas and several other tumors. This MA... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [CEA31] |
| Molekulargewicht | 80-200 kDa |
| UniProt | P06731 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |