

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cdc20(AR12), CF647 conjugate, 0.1mg/mL, Clone: [AR12], Mouse, Monoclonal BOT-BNC470672-500**

Artikelname	Cdc20(AR12), CF647 conjugate, 0.1mg/mL, Clone: [AR12], Mouse, Monoclonal
Artikelnummer	BOT-BNC470672-500
Hersteller Artikelnummer	BNC470672-500
Alternativnummer	BOT-BNC470672-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Urea-denatured His6 Cdc20 human recombinant protein
Konjugation	CF647
Produktbeschreibung	Cyclins, regulatory subunits that associate with kinases, control many of the important steps in cell cycle progression. The Cdc2 protein kinase (p34Cdc2) exhibits protein kinase activity in vitro and exists in a complex with both cyclin B and a prot...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[AR12]

Molekulargewicht	55 kDa
UniProt	<a href="#">Q12834</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user