

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 8(H1 + TS1), CF647 conjugate, 0.1mg/mL, Clone: [H1 TS1], Mouse, Monoclonal BOT-BNC470687-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin 8(H1 + TS1), CF647 conjugate, 0.1mg/mL, Clone: [H1 TS1], Mouse, Monoclonal  |
| Artikelnummer            | BOT-BNC470687-100   |
| Hersteller Artikelnummer | BNC470687-100   |
| Alternativnummer         | BOT-BNC470687-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | FC, IF, IHC, WB   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Cytoskeleton preparation containing cytokeratin 8 (H1), Keratin preparation from a human carcinoma (TS1)  |
| Konjugation              | CF647   |
| Produktbeschreibung      | Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenoc... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [H1 TS1]  |
| Molekulargewicht       | 52.5 kDa  |
| UniProt                | <a href="#">P05787</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |