

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Human Lambda Light Chain(LcN-2 + ICO-106), CF647 conjugate, 0.1mg/mL, Clone: [LcN-2 ICO-106], Mouse, Monoclonal BOT-BNC470735-100

Artikelname	Human Lambda Light Chain(LcN-2 + ICO-106), CF647 conjugate, 0.1mg/mL, Clone: [LcN-2 ICO-106], Mouse, Monoclonal
Artikelnummer	BOT-BNC470735-100
Hersteller Artikelnummer	BNC470735-100
Alternativnummer	BOT-BNC470735-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	WB
Spezies Reaktivität	Human
Immunogen	Purified human IgG (LcN-2 and ICO-106)
Konjugation	CF647
Produktbeschreibung	This MAb is specific to lambda light chain of immunoglobulin and shows no cross-reaction with lambda light chain or any of the five heavy chains. In mammals, the two light chains in an antibody are always identical, with only one type of light chain,...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[LcN-2 ICO-106]

Molekulargewicht	~22.5 kDa
UniProt	P01701
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1.0 ug/mL Optimal dilution for a specific application should be determined by user</p>