

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 17(KRT17/778), CF647 conjugate, 0.1mg/mL, Clone: [KRT17/778], Mouse, Monoclonal BOT-BNC470778-500

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 17(KRT17/778), CF647 conjugate, 0.1mg/mL, Clone: [KRT17/778], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470778-500 |
| Hersteller Artikelnummer | BNC470778-500 |
| Alternativnummer | BOT-BNC470778-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Bovine, Goat, Human, Porcine, Rat |
| Immunogen | Recombinant full-length human KRT17 protein |
| Konjugation | CF647 |
| Produktbeschreibung | Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KRT17/778] |

| | |
|------------------------|--|
| Molekulargewicht | 46 kDa |
| UniProt | Q04695 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |