

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD30(Ki-1/779), CF647 conjugate, 0.1mg/mL, Clone: [Ki-1/779], Mouse, Monoclonal BOT-BNC470779-500**

Artikelname	CD30(Ki-1/779), CF647 conjugate, 0.1mg/mL, Clone: [Ki-1/779], Mouse, Monoclonal
Artikelnummer	BOT-BNC470779-500
Hersteller Artikelnummer	BNC470779-500
Alternativnummer	BOT-BNC470779-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human TNFRSF8 protein
Konjugation	CF647
Produktbeschreibung	Recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In Hodgkins disea...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[Ki-1/779]
Molekulargewicht	105-120 kDa
UniProt	<a href="#">P28908</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user