

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 19(KRT19/800), CF647 conjugate, 0.1mg/mL, Clone: [KRT19/800], Mouse, Monoclonal BOT-BNC470800-100**

Artikelname	Cytokeratin 19(KRT19/800), CF647 conjugate, 0.1mg/mL, Clone: [KRT19/800], Mouse, Monoclonal
Artikelnummer	BOT-BNC470800-100
Hersteller Artikelnummer	BNC470800-100
Alternativnummer	BOT-BNC470800-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IF, WB
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant human KRT19 protein
Konjugation	CF647
Produktbeschreibung	Recognizes a protein of 40 kDa, identified as cytokeratin-19 (CK19), which is expressed in sweat gland, mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, and ectocervical epit...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[KRT19/800]

Molekulargewicht	40 kDa
UniProt	<a href="#">P08727</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user