

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MAGE A1(MZ2E/838), CF647 conjugate, 0.1mg/mL, Clone: [MZ2E/838], Mouse, Monoclonal BOT-BNC470838-100

| | |
|--------------------------|---|
| Artikelname | MAGE A1(MZ2E/838), CF647 conjugate, 0.1mg/mL, Clone: [MZ2E/838], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470838-100 |
| Hersteller Artikelnummer | BNC470838-100 |
| Alternativnummer | BOT-BNC470838-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human MAGEA1 protein |
| Konjugation | CF647 |
| Produktbeschreibung | Recognizes a protein of 42-46 kDa, identified as MAGE-1. This MAb does not cross-react with other members of MAGE-family. Human malignant neoplasms carry rejection antigens that are recognized by the patients autologous, tumor directed and specific, ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [MZ2E/838] |

| | |
|------------------------|--|
| Molekulargewicht | 42-46 kDa |
| UniProt | P43355 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |