

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 10(KRT10/844), CF647 conjugate, 0.1mg/mL, Clone: [KRT10/844], Mouse, Monoclonal BOT-BNC470844-100**

|                          |   |
|--------------------------|---|
| Artikelname              | Cytokeratin 10(KRT10/844), CF647 conjugate, 0.1mg/mL, Clone: [KRT10/844], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC470844-100   |
| Hersteller Artikelnummer | BNC470844-100   |
| Alternativnummer         | BOT-BNC470844-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human, Mouse  |
| Immunogen                | Recombinant human KRT10 protein   |
| Konjugation              | CF647   |
| Produktbeschreibung      | This MAb recognizes a protein of 56.5 kDa, identified as cytokeratin 10 (CK10). CK10 is expressed in all suprabasal layers of the epidermis. In the epidermis, expression of CK10 strictly parallels the extent of differentiation, it is absent in the ba... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [KRT10/844]   |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 56.5 kDa   |
| UniProt                | <a href="#">P13645</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.1-0.2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |