

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HSP60(LK1), CF647 conjugate, 0.1mg/mL, Clone: [LK1], Mouse, Monoclonal BOT-BNC470851-100

| | |
|--------------------------|---|
| Artikelname | HSP60(LK1), CF647 conjugate, 0.1mg/mL, Clone: [LK1], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470851-100 |
| Hersteller Artikelnummer | BNC470851-100 |
| Alternativnummer | BOT-BNC470851-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, IP, WB |
| Spezies Reaktivität | Bovine, Canine, Drosophila, Gallus, Hamster, Human, Monkey, Mouse, Porcine, Rabbit, Rat, Sheep, Xenopus |
| Immunogen | Recombinant human HSP60 protein |
| Konjugation | CF647 |
| Produktbeschreibung | Recognizes a 60 kDa protein, identified as the heat shock protein 60 (hsp60). Its epitope is localized between aa 383-447 of human hsp60. A wide variety of environmental and pathophysiological stressful conditions trigger the synthesis of a family of... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [LK1] |
| Molekulargewicht | 60 kDa |
| UniProt | P10809 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Does not react with bacteria, helminths, or spinach, others not known Western blotting 0.25-0.5 ug/mL Optimal dilution for a specific application should be determined by user</p> |