

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

WT1 (Wilms Tumor 1)(6F-H2), CF647 conjugate, 0.1mg/mL, Clone: [6F-H2], Mouse, Monoclonal BOT-BNC470856-100

Artikelname	WT1 (Wilms Tumor 1)(6F-H2), CF647 conjugate, 0.1mg/mL, Clone: [6F-H2], Mouse, Monoclonal
Artikelnummer	BOT-BNC470856-100
Hersteller Artikelnummer	BNC470856-100
Alternativnummer	BOT-BNC470856-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse, Rat
Immunogen	Recombinant protein corresponding to residues 1-181 of human WT1.
Konjugation	CF647
Produktbeschreibung	Recognizes a 47-55 kDa-tumor suppressor protein, identified as Wilms Tumor (WT1) protein. The antibody reacts with all isoforms of the full-length WT1 and also identifies WT1 lacking exon 2-encoded amino acids, frequently found in subsets of sporadic...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[6F-H2]
Molekulargewicht	47-55 kDa
UniProt	P19544
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user