

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HSP27(G3.1), CF647 conjugate, 0.1mg/mL, Clone: [G3.1], Mouse, Monoclonal BOT-BNC470861-100

| | |
|--------------------------|---|
| Artikelname | HSP27(G3.1), CF647 conjugate, 0.1mg/mL, Clone: [G3.1], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470861-100 |
| Hersteller Artikelnummer | BNC470861-100 |
| Alternativnummer | BOT-BNC470861-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, IP, WB |
| Spezies Reaktivität | Gallus, Human, Monkey, Mouse, Primate, Rat, Sheep |
| Immunogen | Partially purified hsp27 (earlier called 24K) protein from breast cancer MCF-7 cells. |
| Konjugation | CF647 |
| Produktbeschreibung | This MAb reacts specifically with heat shock protein HSP27 in human and monkey tissues and cell lines such as MCF-7. HSP27, also referred to as the Estrogen-Regulated 24K protein and HSP28, is one of several small heat shock proteins produced by all ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [G3.1] |
| Molekulargewicht | 27 kDa |
| UniProt | P04792 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.25-0.5 ug/mL Optimal dilution for a specific application should be determined by user |