

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Glypican-3(GPC3/863 + 1G12), CF647 conjugate, 0.1mg/mL, Clone: [GPC3/863 1G12], Mouse, Monoclonal BOT-BNC470864-100

| | |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname | Glypican-3(GPC3/863 + 1G12), CF647 conjugate, 0.1mg/mL, Clone: [GPC3/863 1G12], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470864-100 |
| Hersteller Artikelnummer | BNC470864-100 |
| Alternativnummer | BOT-BNC470864-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant fragment containing amino acids 511-580 of human glypican-3 (1G12), Recombinant full-length human GPC3 protein (GPC3/863) |
| Konjugation | CF647 |
| Produktbeschreibung | Glypican-3 (GPC3) is an integral membrane protein that is mutated in the Simpson-Golabi-Behmel syndrome (SGBS). SGBS is characterized by pre- and post-natal overgrowth and is a recessive X-linked condition. GPC3 may also be found in a secreted form. ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Klon-Bezeichnung | [GPC3/863 1G12] |
| Molekulargewicht | 67 kDa |
| UniProt | P51654 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |