

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p57 / KIP2(KIP2/880), CF647 conjugate, 0.1mg/mL, Clone: [KIP2/880], Mouse, Monoclonal BOT-BNC470880-100

Artikelname	p57 / KIP2(KIP2/880), CF647 conjugate, 0.1mg/mL, Clone: [KIP2/880], Mouse, Monoclonal
Artikelnummer	BOT-BNC470880-100
Hersteller Artikelnummer	BNC470880-100
Alternativnummer	BOT-BNC470880-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human, Mouse
Immunogen	Recombinant human p57Kip2 protein
Konjugation	CF647
Produktbeschreibung	Recognizes a protein of 57 kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as a...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[KIP2/880]

Molekulargewicht	57 kDa
UniProt	P49918
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user