

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

NGFR (Nerve Growth Factor Receptor)(NGFR5), CF647 conjugate, 0.1mg/mL, Clone: [NGFR5], Mouse, Monoclonal BOT-BNC470890-500

| | |
|--------------------------|---|
| Artikelname | NGFR (Nerve Growth Factor Receptor)(NGFR5), CF647 conjugate, 0.1mg/mL, Clone: [NGFR5], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470890-500 |
| Hersteller Artikelnummer | BNC470890-500 |
| Alternativnummer | BOT-BNC470890-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Feline, Ferret, Human, Monkey, Primate, Rabbit |
| Immunogen | NGFR from A875 melanoma cells |
| Konjugation | CF647 |
| Produktbeschreibung | This antibody recognizes a glycoprotein of 75 kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). Its epitope spans in aa 1-160 of extracellular domain of NGFR/NTR. NGF-receptor contains an ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [NGFR5] |

| | |
|------------------------|---|
| Molekulargewicht | 75 kDa |
| UniProt | P08138 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with mouse or rat, others not known Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |