

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Melanoma Marker(KBA.62), CF647 conjugate, 0.1mg/mL, Clone: [KBA.62], Mouse, Monoclonal BOT-BNC470895-500

| | |
|--------------------------|---|
| Artikelname | Melanoma Marker(KBA.62), CF647 conjugate, 0.1mg/mL, Clone: [KBA.62], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470895-500 |
| Hersteller Artikelnummer | BNC470895-500 |
| Alternativnummer | BOT-BNC470895-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | Human KAL cells derived from lymph node metastasis of malignant melanoma |
| Konjugation | CF647 |
| Produktbeschreibung | KBA.62 is a novel anti-melanoma antibody. It reacts positively against melanocytic tumors but not other tumors, thus demonstrating specificity and sensitivity. Moreover, it reacts positively against junctional nevus cells but not intradermal nevi, an... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|---|
| Klon-Bezeichnung | [KBA.62] |
| Molekulargewicht | Multiple (140, 135 and 128 kDa and two weak bands of 88 and 73 kDa) |
| UniProt | Not Known |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user |