

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

MiTF (Microphthalmia Transcription Factor)(D5 + MITF/915), CF647 conjugate, 0.1mg/mL, Clone: [D5 MITF/915], Mouse, Monoclonal BOT-BNC470950-500

| | |
|--------------------------|---|
| Artikelname | MiTF (Microphthalmia Transcription Factor)(D5 + MITF/915), CF647 conjugate, 0.1mg/mL, Clone: [D5 MITF/915], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470950-500 |
| Hersteller Artikelnummer | BNC470950-500 |
| Alternativnummer | BOT-BNC470950-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human |
| Immunogen | NH2 terminus fragment of human Mi protein (D5), Recombinant human MiTF protein (MITF/915) |
| Konjugation | CF647 |
| Produktbeschreibung | MITF (microphthalmia transcription factor) is a basic helix-loop-helix-leucine-zipper (bHLH-Zip) transcription factor that regulates the development and survival of melanocytes and retinal pigment epithelium, and also is involved in transcription of ... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [D5 MITF/915] |
| Molekulargewicht | 52-56 kDa (doublet) |
| UniProt | O75030 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with mouse or rat, others not tested Immunohistology (formalin) Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |