

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Myeloid-Associated Differentiation Marker(MYADM/971), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [MYADM/971], Mouse, Monoclonal BOT-BNC470971-500

| | |
|--------------------------|---|
| Artikelname | Myeloid-Associated Differentiation Marker(MYADM/971), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [MYADM/971], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470971-500 |
| Hersteller Artikelnummer | BNC470971-500 |
| Alternativnummer | BOT-BNC470971-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC |
| Spezies Reaktivität | Human, Primate |
| Immunogen | Recombinant human MYADM protein |
| Konjugation | CF647 |
| Produktbeschreibung | Recognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellul... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [MYADM/971] |

| | |
|------------------------|---|
| Molekulargewicht | Unknown |
| Isotyp | IgG1 |
| UniProt | Q96S97 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |