

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TGF-beta(1D11.16.8), CF647 conjugate, 0.1mg/mL, Clone: [1D11.16.8], Mouse, Monoclonal BOT-BNC470977-500

| | |
|--------------------------|---|
| Artikelname | TGF-beta(1D11.16.8), CF647 conjugate, 0.1mg/mL, Clone: [1D11.16.8], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC470977-500 |
| Hersteller Artikelnummer | BNC470977-500 |
| Alternativnummer | BOT-BNC470977-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | Functional Studies, IHC |
| Spezies Reaktivität | Bovine, Canine, Hamster, Human, Monkey, Mouse, Rat |
| Immunogen | Bovine TGF-beta2 protein |
| Konjugation | CF647 |
| Produktbeschreibung | This MAb recognizes TGF beta 1, 2 and 3. Three TGF-betas have been identified in mammals. TGF-beta1, TGF-beta2 and TGF-beta3 are each synthesized as precursor proteins that are very similar in that each is cleaved to yield a 112 amino acid polypeptid... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [1D11.16.8] |

| | |
|------------------------|--|
| Molekulargewicht | ~13 kDa |
| UniProt | P01137 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Cytokine neutralization or functional assay, order Ab without BSA/azide Optimal dilution for a specific application should be determined by user |