

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

p57 / KIP2(KP10 + KIP2/880), CF647 conjugate, 0.1mg/mL, Clone: [KP10 KIP2/880], Mouse, Monoclonal BOT-BNC471015-100

| | |
|--------------------------|---|
| Artikelname | p57 / KIP2(KP10 + KIP2/880), CF647 conjugate, 0.1mg/mL, Clone: [KP10 KIP2/880], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471015-100 |
| Hersteller Artikelnummer | BNC471015-100 |
| Alternativnummer | BOT-BNC471015-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human, Mouse |
| Immunogen | Recombinant human p57Kip2 protein |
| Konjugation | CF647 |
| Produktbeschreibung | Recognizes a protein of 57 kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as a... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [KP10 KIP2/880] |
| Molekulargewicht | 57 kDa |

| | |
|------------------------|--|
| UniProt | P49918 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |