

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Beta-2-Microglobulin (B2M)(246-E9.E7), CF647 conjugate, 0.1mg/mL, Clone: [246-E9.E7|same as HLA.ABC.m2], Mouse, Monoclonal BOT-BNC471039-100

| | |
|--------------------------|---|
| Artikelname | Beta-2-Microglobulin (B2M)(246-E9.E7), CF647 conjugate, 0.1mg/mL, Clone: [246-E9.E7 same as HLA.ABC.m2], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471039-100 |
| Hersteller Artikelnummer | BNC471039-100 |
| Alternativnummer | BOT-BNC471039-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, Functional Studies, IF, IP |
| Spezies Reaktivität | Human |
| Immunogen | Human PBLs from a T-cell acute lymphoblastic leukemia (T-ALL) patient |
| Konjugation | CF647 |
| Produktbeschreibung | Beta-2 microglobulin is a component of the class I major histocompatibility complex (MHC I) where it functions in antigen processing and presentation. Involved in the presentation of peptide antigens to the immune system. Human Beta-2 microglobulin i... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [246-E9.E7 same as HLA.ABC.m2] |
| Molekulargewicht | 12 kDa |
| UniProt | P61769 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Does not react with non-human primates, others not known Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |