

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

HLA-DRB(HLA-DRB/1067), CF647 conjugate, 0.1mg/mL, Clone: [HLA-DRB/1067], Mouse, Monoclonal BOT-BNC471067-100

| | |
|--------------------------|---|
| Artikelname | HLA-DRB(HLA-DRB/1067), CF647 conjugate, 0.1mg/mL, Clone: [HLA-DRB/1067], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471067-100 |
| Hersteller Artikelnummer | BNC471067-100 |
| Alternativnummer | BOT-BNC471067-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human, Monkey |
| Immunogen | Activated human peripheral blood mononuclear cells |
| Konjugation | CF647 |
| Produktbeschreibung | This MAb reacts with the beta-chain of HLA-DRB1 antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. Its epitope is different from that of MAb L243. HLA-DR is a heterodimeric cell surface glycoprotein comprised... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [HLA-DRB/1067] |

| | |
|------------------------|--|
| Molekulargewicht | 28 kDa (beta chain) |
| UniProt | P01911 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.25-0.5 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |