

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ksp-Cadherin / CDH16(CDH16/1071), CF647 conjugate, 0.1mg/mL, Clone: [CDH16/1071], Mouse, Monoclonal BOT-BNC471071-500

| | |
|--------------------------|---|
| Artikelname | Ksp-Cadherin / CDH16(CDH16/1071), CF647 conjugate, 0.1mg/mL, Clone: [CDH16/1071], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471071-500 |
| Hersteller Artikelnummer | BNC471071-500 |
| Alternativnummer | BOT-BNC471071-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IF, IHC, WB |
| Spezies Reaktivität | Canine, Human, Mouse, Rabbit |
| Immunogen | Recombinant human CDH16 protein |
| Konjugation | CF647 |
| Produktbeschreibung | This MAb recognizes a protein of 130 kDa, identified as Ksp-cadherin. Cadherins form a superfamily of related glycoproteins that mediate calcium-dependent cell adhesion and transmit signals from the extracellular matrix to the cytoplasm. Cadherins ha... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [CDH16/1071] |

| | |
|------------------------|---|
| Molekulargewicht | 130 kDa |
| UniProt | O75309 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |