

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MHC II (HLA-DRA)(19-26.1), CF647 conjugate, 0.1mg/mL, Clone: [19-26.1], Mouse, Monoclonal BOT-BNC471082-100**

Artikelname	MHC II (HLA-DRA)(19-26.1), CF647 conjugate, 0.1mg/mL, Clone: [19-26.1], Mouse, Monoclonal
Artikelnummer	BOT-BNC471082-100
Hersteller Artikelnummer	BNC471082-100
Alternativnummer	BOT-BNC471082-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IHC
Spezies Reaktivität	Human
Immunogen	Raji cells
Konjugation	CF647
Produktbeschreibung	This MAb reacts with the HLA-DR antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kDa alpha (heavy) chain and a 28 kDa beta (light) chain...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[19-26.1]

Molekulargewicht	36 kDa (? chain) and 27 kDa (? chain)
UniProt	<a href="#">P01903</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 1-2 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user