

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TSH-Receptor, B-Chain (Thyroid Marker) (TSHRB/1404), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [TSHRB/1404], Mouse, Monoclonal BOT-BNC471404-500

| | |
|--------------------------|---|
| Artikelname | TSH-Receptor, B-Chain (Thyroid Marker) (TSHRB/1404), CF647 conjugate, 0.1mg/mL, IgG1, Clone: [TSHRB/1404], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471404-500 |
| Hersteller Artikelnummer | BNC471404-500 |
| Alternativnummer | BOT-BNC471404-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | B-Chain protein Recombinant human TSHR |
| Konjugation | CF647 |
| Produktbeschreibung | Thyroid-stimulating hormone (TSH, also known as thyrotropin) is a glycoprotein involved in the control of thyroid structure and metabolism, which stimulates the release of the thyroid hormones. TSH is regulated by thyroid hormone (T3) and various ret... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [TSHRB/1404] |

| | |
|------------------------|--|
| Molekulargewicht | 42 kDa |
| Isotyp | IgG1 |
| UniProt | P16473 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |