

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker), CF647 conjugate, 0.1mg/mL, Rabbit BOT-BNC471536-100**

|                          |                                                                                                                                                                                                                                                               |
|--------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Artikelname              | CD30 / TNFRSF8 (Hodgkin & Reed-Sternberg Cell Marker), CF647 conjugate, 0.1mg/mL, Rabbit                                                                                                                                                                      |
| Artikelnummer            | BOT-BNC471536-100                                                                                                                                                                                                                                             |
| Hersteller Artikelnummer | BNC471536-100                                                                                                                                                                                                                                                 |
| Alternativnummer         | BOT-BNC471536-100-100UL                                                                                                                                                                                                                                       |
| Hersteller               | Biotium                                                                                                                                                                                                                                                       |
| Wirt                     | Rabbit                                                                                                                                                                                                                                                        |
| Kategorie                | Antikörper                                                                                                                                                                                                                                                    |
| Applikation              | IHC                                                                                                                                                                                                                                                           |
| Spezies Reaktivität      | Human                                                                                                                                                                                                                                                         |
| Immunogen                | Recombinant human full-length TNFRSF8 protein                                                                                                                                                                                                                 |
| Konjugation              | CF647                                                                                                                                                                                                                                                         |
| Produktbeschreibung      | This antibody recognizes a single chain glycoprotein of 105/120 kDa, identified as CD30/Ki-1. CD30 is synthesized as a 90 kDa precursor, which is processed in the Golgi complex into a membrane-bound phosphorylated mature 105/120 kDa glycoprotein. In ... |
| Konzentration            | 0.1 mg/mL                                                                                                                                                                                                                                                     |
| Molekulargewicht         | 105-120 kDa                                                                                                                                                                                                                                                   |
| UniProt                  | <a href="#">P28908</a>                                                                                                                                                                                                                                        |

|                        |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Puffer                 | PBS, 0.1% BSA, 0.05% azide                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Quelle                 | Animal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA pH 8.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p> |