

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Filaggrin (Keratinocyte Differentiation Marker) (FLG/1562), CF647 conjugate, 0.1mg/mL, Clone: [FLG/1562], Mouse, Monoclonal BOT-BNC471562-500

Artikelname	Filaggrin (Keratinocyte Differentiation Marker) (FLG/1562), CF647 conjugate, 0.1mg/mL, Clone: [FLG/1562], Mouse, Monoclonal
Artikelnummer	BOT-BNC471562-500
Hersteller Artikelnummer	BNC471562-500
Alternativnummer	BOT-BNC471562-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human Filaggrin protein fragment (aa 998-1104) (exact sequence is proprietary)
Konjugation	CF647
Produktbeschreibung	Filaggrin is an intermediate filament-associated protein that aggregates keratin intermediate filaments in mammalian epidermis. It is initially synthesized as a polyprotein precursor, profilaggrin (consisting of multiple filaggrin units of 324 aa eac...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[FLG/1562]
Molekulargewicht	26-45 kDa (Processed), 350 kDa (Profilaggrin)
UniProt	P20930
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user