

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **SREBP1(SREBP1/1578), CF647 conjugate, 0.1mg/mL, Clone: [SREBP1/1578], Mouse, Monoclonal BOT-BNC471578-100**

|                          |   |
|--------------------------|---|
| Artikelname              | SREBP1(SREBP1/1578), CF647 conjugate, 0.1mg/mL, Clone: [SREBP1/1578], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC471578-100   |
| Hersteller Artikelnummer | BNC471578-100   |
| Alternativnummer         | BOT-BNC471578-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Spezies Reaktivität      | Human   |
| Immunogen                | Recombinant full-length human SREBF1 protein  |
| Konjugation              | CF647   |
| Produktbeschreibung      | The low density lipoprotein (LDL) receptor mediates the endocytic uptake of cholesterol-carrying lipoproteins, thereby controlling cholesterol levels in cells and plasma. Transcription of the LDL receptor gene is controlled by a ten base pair sequenc... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [SREBP1/1578]   |
| Molekulargewicht         | 68/125 kDa mature/precursor   |

|                        |  |
|------------------------|--|
| UniProt                | <a href="#">P36956</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |