

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD80 (B7-1) (C80/1608), CF647 conjugate, 0.1mg/mL, Clone: [C80/1608], Mouse, Monoclonal BOT-BNC471608-500

| | |
|--------------------------|---|
| Artikelname | CD80 (B7-1) (C80/1608), CF647 conjugate, 0.1mg/mL, Clone: [C80/1608], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC471608-500 |
| Hersteller Artikelnummer | BNC471608-500 |
| Alternativnummer | BOT-BNC471608-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human full-length CD80 protein |
| Konjugation | CF647 |
| Produktbeschreibung | T cell proliferation and lymphokine production are triggered by occupation of the TCR by antigen, followed by a costimulatory signal that is delivered by a ligand expressed on antigen presenting cells. The B7-related cell surface proteins CD80 (B7-1)... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [C80/1608] |
| Molekulargewicht | 60 kDa |

| | |
|------------------------|--|
| UniProt | P33681 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |