

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Fibronectin (8th Type III Repeat)(568), CF568 conjugate, 0.1mg/mL, Clone: [568], Mouse, Monoclonal BOT-BNC680351-100

Artikelname	Fibronectin (8th Type III Repeat)(568), CF568 conjugate, 0.1mg/mL, Clone: [568], Mouse, Monoclonal
Artikelnummer	BOT-BNC680351-100
Hersteller Artikelnummer	BNC680351-100
Alternativnummer	BOT-BNC680351-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	High molecular weight proteins secreted by cultivated human fibroblasts
Konjugation	CF568
Produktbeschreibung	The epitope of this antibody is located in the 8th type III repeat in the cell-binding region of fibronectin. Fibronectins are disulfide-linked, dimeric glycoproteins of ~440 kDa. They possess at least four binding sites for collagen, glycosaminoglyc...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[568]
Molekulargewicht	220 kDa (monomer), 440 kDa (dimer)
UniProt	P02751
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistochemistry (formalin-fixed): 0.5-1 ug/mL for 30 minutes at RT Flow cytometry: 0.5-1 ug/million cells Immunofluorescence: 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris buffer with 1 mM EDTA pH 9.0 for 10-20 minutes followed by cooling at RT for 20 minutes Optimal dilution for a specific application should be determined by user