

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Human IgM Immunoglobulin(ICO-30), CF568 conjugate, 0.1mg/mL, Clone: [ICO-30], Mouse, Monoclonal BOT-BNC680364-500**

|                          |   |
|--------------------------|---|
| Artikelname              | Human IgM Immunoglobulin(ICO-30), CF568 conjugate, 0.1mg/mL, Clone: [ICO-30], Mouse, Monoclonal   |
| Artikelnummer            | BOT-BNC680364-500   |
| Hersteller Artikelnummer | BNC680364-500   |
| Alternativnummer         | BOT-BNC680364-500-500UL   |
| Hersteller               | Biotium   |
| Wirt                     | Mouse   |
| Kategorie                | Antikörper  |
| Applikation              | IHC   |
| Spezies Reaktivität      | Human   |
| Immunogen                | Heavy chain of human IgM  |
| Konjugation              | CF568   |
| Produktbeschreibung      | Recognizes a protein of 75 kDa, identified as mu heavy chain of human immunoglobulins. It does not cross-react with alpha (IgA), gamma (IgG), epsilon (IgE), or delta (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. Monomeric Ig... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |
| Klon-Bezeichnung         | [ICO-30]  |

|                        |  |
|------------------------|--|
| Molekulargewicht       | 50-75 kDa  |
| UniProt                | <a href="#">P01871</a>   |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide   |
| Quelle                 | Animal   |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |