

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **CD81 / TAPA-1 (1.3.3.22), CF568 conjugate, 0.1mg/mL, Clone: [1.3.3.22], Mouse, Monoclonal BOT-BNC680391-500**

Artikelname	CD81 / TAPA-1 (1.3.3.22), CF568 conjugate, 0.1mg/mL, Clone: [1.3.3.22], Mouse, Monoclonal
Artikelnummer	BOT-BNC680391-500
Hersteller Artikelnummer	BNC680391-500
Alternativnummer	BOT-BNC680391-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	Exosome Staining, FC, Functional Studies, IF, IHC, WB
Spezies Reaktivität	Human, Rat
Immunogen	B-Cell line derived from a Burkitt lymphoma
Konjugation	CF568
Produktbeschreibung	This antibody recognizes a protein of 26 kDa, identified as CD81 (Workshop VI, Code CD81.1). CD81 has a very broad cellular distribution, being expressed on T- and B-lymphocytes, NK cells, thymocytes, eosinophils, fibroblasts, epithelial and endothel...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[1.3.3.22]

Molekulargewicht	26 kDa
UniProt	<a href="#">P60033</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL For functional studies, order Ab without azide Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user