

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Cytokeratin 14(LL002), CF568 conjugate, 0.1mg/mL, IgG3, Clone: [LL002], Mouse, Monoclonal BOT-BNC680499-500

| | |
|--------------------------|---|
| Artikelname | Cytokeratin 14(LL002), CF568 conjugate, 0.1mg/mL, IgG3, Clone: [LL002], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC680499-500 |
| Hersteller Artikelnummer | BNC680499-500 |
| Alternativnummer | BOT-BNC680499-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC |
| Spezies Reaktivität | Human, Mouse, Rat |
| Immunogen | A synthetic peptide of 15 amino acid from the C-terminus of human keratin 14. |
| Konjugation | CF568 |
| Produktbeschreibung | Cytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B or basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, my... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [LL002] |
| Molekulargewicht | ~50 kDa |
| Isotyp | IgG3 |
| UniProt | P02533 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology formalin-fixed 0.5-1.0 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user |