

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

TGF-beta (Transforming Growth Factor beta)(TGFB/510), CF568 conjugate, 0.1mg/mL, Clone: [TGFB/510], Mouse, Monoclonal BOT-BNC680510-100

| | |
|--------------------------|---|
| Artikelname | TGF-beta (Transforming Growth Factor beta)(TGFB/510), CF568 conjugate, 0.1mg/mL, Clone: [TGFB/510], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC680510-100 |
| Hersteller Artikelnummer | BNC680510-100 |
| Alternativnummer | BOT-BNC680510-100-100UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Spezies Reaktivität | Bovine, Canine, Hamster, Human, Monkey, Mouse, Rat |
| Immunogen | Full length recombinant human TGF-beta2 protein |
| Konjugation | CF568 |
| Produktbeschreibung | This MAb recognizes TGF beta 1, 2 and 3. TGFs inhibit the growth of epithelial cells and stimulate the growth of mesenchymal cells. Most breast lesions, benign and malignant, involve abnormal proliferation and altered architecture of stromal and/or e... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [TGFB/510] |
| Molekulargewicht | ~13 kDa |

| | |
|------------------------|--|
| UniProt | P01137 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | For coating for ELISA, order Ab without BSA Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Optimal dilution and staining procedure for a specific application should be determined by user Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry |