

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Cytokeratin 18(DE-K18), CF568 conjugate, 0.1mg/mL, Clone: [DE-K18], Mouse, Monoclonal BOT-BNC680563-100**

Artikelname	Cytokeratin 18(DE-K18), CF568 conjugate, 0.1mg/mL, Clone: [DE-K18], Mouse, Monoclonal
Artikelnummer	BOT-BNC680563-100
Hersteller Artikelnummer	BNC680563-100
Alternativnummer	BOT-BNC680563-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	FC, IF, IHC, WB
Spezies Reaktivität	Canine, Feline, Human
Immunogen	Cytoskeletal preparation extracted from the human vulvar squamous carcinoma cell line A431
Konjugation	CF568
Produktbeschreibung	This MAb reacts with a wide variety of simple epithelia. It does not react with stratified squamous epithelia. It reacts with epithelial tumors of the gastrointestinal tract, lung, breast, pancreas, ovary, and thyroid. Cytokeratin 18, which belongs t...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[DE-K18]
Molekulargewicht	45 kDa
UniProt	<a href="#">P05783</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	<p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Does not react with rat, others not known Immunohistology formalin-fixed 2-4 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user</p>