

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **TDP2 / EAPII (TDP2/1258), CF568 conjugate, 0.1mg/mL, Clone: [TDP2/1258], Mouse, Monoclonal BOT-BNC681258-500**

Artikelname	TDP2 / EAPII (TDP2/1258), CF568 conjugate, 0.1mg/mL, Clone: [TDP2/1258], Mouse, Monoclonal
Artikelnummer	BOT-BNC681258-500
Hersteller Artikelnummer	BNC681258-500
Alternativnummer	BOT-BNC681258-500-500UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human full-length TDP2 protein
Konjugation	CF568
Produktbeschreibung	This MAb recognizes a protein of 41 kDa, which is identified as TDP2. It is a member of a superfamily of divalent cation-dependent phosphodiesterases. The encoded protein associates with CD40, tumor necrosis factor (TNF) receptor-75 and TNF receptor ...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[TDP2/1258]

Molekulargewicht	~41 kDa
UniProt	<a href="#">O95551</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user