

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

CD3e (T-Cell Marker) (C3e/1308), CF568 conjugate, 0.1mg/mL, Clone: [C3e/1308], Mouse, Monoclonal BOT-BNC681308-500

| | |
|--------------------------|---|
| Artikelname | CD3e (T-Cell Marker) (C3e/1308), CF568 conjugate, 0.1mg/mL, Clone: [C3e/1308], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681308-500 |
| Hersteller Artikelnummer | BNC681308-500 |
| Alternativnummer | BOT-BNC681308-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human CD3e extracellular domain fragment (aa 23-119) |
| Konjugation | CF568 |
| Produktbeschreibung | The immunogen for this Mab is derived from the extracellular domain of CD3 epsilon chain. CD3 consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closel... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [C3e/1308] |
| Molekulargewicht | 20 kDa |
| UniProt | P07766 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Not recommended for flow cytometry Immunohistology (formalin) 1-2 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Western blotting 1-2 ug/mL Immunofluorescence 1-2 ug/mL Optimal dilution for a specific application should be determined by user |