

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Protocadherin FAT2 (FAT Atypical Cadherin 2) (8C5), CF568 conjugate, 0.1mg/mL, Clone: [8C5], Mouse, Monoclonal BOT-BNC681320-100

Artikelname	Protocadherin FAT2 (FAT Atypical Cadherin 2) (8C5), CF568 conjugate, 0.1mg/mL, Clone: [8C5], Mouse, Monoclonal
Artikelnummer	BOT-BNC681320-100
Hersteller Artikelnummer	BNC681320-100
Alternativnummer	BOT-BNC681320-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Purified recombinant human FAT2 fusion protein
Konjugation	CF568
Produktbeschreibung	This antibody recognizes a protein of 480 kDa, which is identified as FAT2. The cadherins represent a family of Ca ²⁺ -dependent adhesion molecules that function to mediate cell-to-cell binding that is critical for the maintenance of structure and morp...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL
Klon-Bezeichnung	[8C5]

Molekulargewicht	480 kDa
UniProt	Q9NYQ8
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Optimal dilution for a specific application should be determined by user