

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **Adipophilin / Perilipin-2 (Marker of Obesity) (ADFP/1366), CF568 conjugate, 0.1mg/mL, Clone: [ADFP/1366], Mouse, Monoclonal BOT-BNC681366-100**

Artikelname	Adipophilin / Perilipin-2 (Marker of Obesity) (ADFP/1366), CF568 conjugate, 0.1mg/mL, Clone: [ADFP/1366], Mouse, Monoclonal
Artikelnummer	BOT-BNC681366-100
Hersteller Artikelnummer	BNC681366-100
Alternativnummer	BOT-BNC681366-100-100UL
Hersteller	Biotium
Wirt	Mouse
Kategorie	Antikörper
Applikation	IHC
Spezies Reaktivität	Human
Immunogen	Recombinant human Adipophilin protein fragment (aa249-376) (exact sequence is proprietary)
Konjugation	CF568
Produktbeschreibung	Perilipin-2, also known as Adipophilin, belongs to the perilipin family, members of which coat intracellular lipid storage droplets. This 48 kDa protein is associated with the lipid globule surface membrane material, and maybe involved in development...
Klonalität	Monoclonal
Konzentration	0.1 mg/mL

Klon-Bezeichnung	[ADFP/1366]
Molekulargewicht	48 kDa
UniProt	<a href="#">Q99541</a>
Puffer	PBS, 0.1% BSA, 0.05% azide
Quelle	Animal
Anwendungsbeschreibung	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min ELISA For coating, order Ab without BSA Optimal dilution for a specific application should be determined by user