

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) (EGP40/1372), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [EGP40/1372], Mouse, Monoclonal BOT-BNC681372-500

| | |
|--------------------------|---|
| Artikelname | Ep-CAM / CD326 (Extracellular Domain) (Epithelial Marker) (EGP40/1372), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [EGP40/1372], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681372-500 |
| Hersteller Artikelnummer | BNC681372-500 |
| Alternativnummer | BOT-BNC681372-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | FC, IF, IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human EpCAM protein fragment from the extracellular domain (aa77-202) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | EGP40 is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majo... |
| Klonalität | Monoclonal |

| | |
|------------------------|--|
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [EGP40/1372] |
| Molekulargewicht | 40-43 kDa |
| Isotyp | IgG1 |
| UniProt | P16422 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |