

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

ALDH1A1 (Aldehyde Dehydrogenase 1A1) (ALDH1A1/1381), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [ALDH1A1/1381], Mouse, Monoclonal BOT-BNC681381-500

| | |
|--------------------------|---|
| Artikelname | ALDH1A1 (Aldehyde Dehydrogenase 1A1) (ALDH1A1/1381), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [ALDH1A1/1381], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681381-500 |
| Hersteller Artikelnummer | BNC681381-500 |
| Alternativnummer | BOT-BNC681381-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | IHC, WB |
| Spezies Reaktivität | Human |
| Immunogen | Purified recombinant fragment of human ALDH1A1 (aa 315-434) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | Aldehyde dehydrogenase 1 family member A1 (ALDH1A1), also known as retinal dehydrogenase 1, belongs to the aldehyde dehydrogenase enzyme family, which is involved in the metabolism of alcohol. ALDH1A1 is predominantly expressed in the epithelium of t... |
| Klonalität | Monoclonal |

| | |
|------------------------|--|
| Konzentration | 0.1 mg/mL |
| Klon-Bezeichnung | [ALDH1A1/1381] |
| Molekulargewicht | 56 kDa |
| Isotyp | IgG1 |
| UniProt | P00352 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 1-2 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user |