

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

Product Datasheet

Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/1389), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [FTL/1389], Mouse, Monoclonal BOT-BNC681389-500

| | |
|--------------------------|---|
| Artikelname | Ferritin, Light Chain (FTL) (Microglia Marker) (FTL/1389), CF568 conjugate, 0.1mg/mL, IgG1, Clone: [FTL/1389], Mouse, Monoclonal |
| Artikelnummer | BOT-BNC681389-500 |
| Hersteller Artikelnummer | BNC681389-500 |
| Alternativnummer | BOT-BNC681389-500-500UL |
| Hersteller | Biotium |
| Wirt | Mouse |
| Kategorie | Antikörper |
| Applikation | WB |
| Spezies Reaktivität | Human |
| Immunogen | Recombinant human FTL protein fragment (aa 38-165) (exact sequence is proprietary) |
| Konjugation | CF568 |
| Produktbeschreibung | Mammalian ferritins consist of 24 subunits made up of 2 types of polypeptide chains, ferritin heavy chain and ferritin light chain. Ferritin heavy chains catalyze the first step in iron storage, the oxidation of Fe (II), whereas ferritin light chains... |
| Klonalität | Monoclonal |
| Konzentration | 0.1 mg/mL |

| | |
|------------------------|--|
| Klon-Bezeichnung | [FTL/1389] |
| Molekulargewicht | 19-25 kDa |
| Isotyp | IgG1 |
| UniProt | P02792 |
| Puffer | PBS, 0.1% BSA, 0.05% azide |
| Quelle | Animal |
| Anwendungsbeschreibung | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunohistology (formalin) 0.1-0.2 ug/mL Flow cytometry 0.1-0.2ug/million cells Immunofluorescence 0.1-0.2ug/ml Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min Optimal dilution for a specific application should be determined by user |