

Bitte beachten Sie: Dieses Dokument wurde automatisch erstellt und ist kein Ersatz für das Originaldokument des Herstellers.

## Product Datasheet

### **MART-1 / Melan-A / MLANA (Melanoma Marker) (MLANA/1409R), CF568 conjugate, 0.1mg/mL, Clone: [MLANA/1409R], Rabbit, Monoclonal BOT-BNC681409-100**

|                          |   |
|--------------------------|---|
| Artikelname              | MART-1 / Melan-A / MLANA (Melanoma Marker) (MLANA/1409R), CF568 conjugate, 0.1mg/mL, Clone: [MLANA/1409R], Rabbit, Monoclonal   |
| Artikelnummer            | BOT-BNC681409-100   |
| Hersteller Artikelnummer | BNC681409-100   |
| Alternativnummer         | BOT-BNC681409-100-100UL   |
| Hersteller               | Biotium   |
| Wirt                     | Rabbit  |
| Kategorie                | Antikörper  |
| Applikation              | IHC, WB   |
| Spezies Reaktivität      | Human, Mouse, Rat   |
| Immunogen                | Recombinant human full-length MLANA protein   |
| Konjugation              | CF568   |
| Produktbeschreibung      | This antibody recognizes a protein doublet of 20-22 kDa, identified as MART-1 (Melanoma Antigen Recognized by T cells 1) or Melan-A. MART-1 is a newly identified melanocyte differentiation antigen recognized by autologous cytotoxic T lymphocytes. Sev... |
| Klonalität               | Monoclonal  |
| Konzentration            | 0.1 mg/mL   |

|                        |   |
|------------------------|---|
| Klon-Bezeichnung       | [MLANA/1409R]   |
| Molekulargewicht       | 20-22 kDa (doublet)   |
| UniProt                | <a href="#">Q16655</a>  |
| Puffer                 | PBS, 0.1% BSA, 0.05% azide  |
| Quelle                 | Animal  |
| Anwendungsbeschreibung | <p>Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody Immunofluorescence: 0.5-1 ug/mL Immunohistology (formalin): 0.5-1 ug/mL Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer pH 6.0 for 10-20 min followed by cooling at RT for 20 min Flow Cytometry 0.5-1 ug/million cells/0.1 mL Western blotting 0.5-1 ug/mL Optimal dilution for a specific application should be determined by user</p> |